



**IDEAL GAS EQUATION NOTES**

$P V = n R T$

"P" stands for \_\_\_\_\_, must be in units of \_\_\_\_\_

"V" stands for \_\_\_\_\_, must be in units of \_\_\_\_\_

"n" stands for \_\_\_\_\_, must be in units of \_\_\_\_\_

"T" stands for \_\_\_\_\_, must be in units of \_\_\_\_\_

"R" stands for the Ideal Gas Constant, has a value of 0.0821 with units of  $\frac{\text{L} \cdot \text{atm}}{\text{mole} \cdot \text{K}}$

**EXAMPLE 1:**

Q: What pressure is exerted by 0.325 moles of hydrogen gas in a 4.08 L container at 35 °C?

A:  $P V = n R T$  solving for P ...  $P = \frac{n R T}{V}$

n = 0.325 moles (correct unit)

T = 35 °C (need to convert to K)  $T = 35 \text{ °C} + 273 = 308 \text{ K}$

V = 4.08 L (correct unit)

$P = \frac{(0.325 \text{ moles}) (0.0821 \frac{\text{L} \cdot \text{atm}}{\text{mole} \cdot \text{K}}) (308 \text{ K})}{(4.08 \text{ L})} = 2.01 \text{ atm}$

**EXAMPLE 2:**

Q: What mass of chlorine gas, Cl<sub>2</sub>, in grams, is contained in a 10.0 L tank at 27 °C and 3.50 atm of pressure? (Answer: 101 grams)

**EXAMPLE 3:**

Q: A gas at 20.0 °C and 3.98 atm contains 1.45 moles of gas particles. What volume does the gas occupy? (Answer: 8.77 L)

**Ideal Gas Equation 1 WKSHT**

- 1.) What is the pressure exerted by 2.0 moles of an ideal gas when it occupies a volume of 12.0 L at 373 K?
- 2.) A flashbulb of volume 2.6 cm<sup>3</sup> contains O<sub>2</sub> gas at a pressure of 2.3 atm and a temperature of 26°C. How many moles of O<sub>2</sub> does the flashbulb contain?
- 3.) If 0.20 moles of helium occupies a volume of 64.0 liters at a pressure of 0.15 atm, what is the temperature of the gas?
- 4.) What is the volume of 0.35 moles of gas at 1.7 atm of pressure and a temperature of 100 K?
- 5.) What is the pressure of 1.5 moles of an ideal gas at a temperature of 150 K and occupies a volume of 20.0 liters?
- 6.) How many moles of gas occupy 16.2 liters at a pressure of 1.05 atm and a temperature of 37°C?

Answers: 1.) 5.1 atm; 2.)  $2.4 \times 10^{-4}$  moles; 3.) 580 K; 4.) 1.7 L; 5.) 0.92 atm; 6.) 0.668 moles

**Ideal Gas Equation 2 WKSHT**

- 1.) Calculate the volume of exactly 1.00 mole of a gas at STP.
- 2.) How many moles of nitrogen are present in 17.8 liters at 27 °C and 1.3 atm pressure?
- 3.) What is the pressure of 2.3 moles of carbon dioxide at 235 K occupying 23.7 liters of space?
- 4.) If there are  $4.02 \times 10^{23}$  molecules of N<sub>2</sub>O in a sample, how many moles are there?

- 5.) Using answer from #6, calculate the pressure of the gas if it occupies 27,025 cm<sup>3</sup> of space at 38.0 °C.
- 6.) How many grams of NH<sub>3</sub> are present in 35.0 dm<sup>3</sup> of space at 78.3 K and 0.853 atm of pressure?
- 7.) What is the temperature of 34.2 grams of sulfur dioxide occupying 30.0 liters of space and having a pressure of 800. torr?
- 8.) What is the pressure (in mm Hg) of 79.4 grams of boron trifluoride in a 20.0 L container at a temperature of 245 K?
- 9.) How many grams are in a sample of arsenic trifluoride that has a volume of 17,600 mL and a temperature of 92 °C and a pressure of 108,732 Pa?
- 10.) How many kilopascals of pressure are exerted by 23.8 liters of oxygen with a mass of 175 grams at a temperature of 58 °C?
- 11.) How many moles of argon are in 30.6 liters at 28 K and 658 mm Hg of pressure?
- 12.) How many grams of argon are found in # 11?

Answers: 1.) 22.4 L; 2.) 0.94 moles; 3.) 1.9 atm; 4.) 0.668 moles; 5.) 0.631 atm; 6.) 78.9 g; 7.) 720. K; 8.) 897 mm Hg; 9.) 83 g; 10.) 632 kPa; 11.) 12 moles; 12.) 480 g

**APPLICATIONS OF THE IDEAL GAS EQUATION NOTES**

EXAMPLE #1:

Q: At 28 °C and 0.974 atm, 1.00 L of gas has a mass of 5.16 grams. What is the molar mass of this gas?

A:  $PV = nRT$       We know moles =  $\frac{\text{grams}}{\text{MM}}$       Using substitution...       $PV = \frac{g}{\text{MM}}RT$

$$\text{MM} \cdot PV = \frac{g}{\text{MM}}RT \cdot \text{MM} \qquad \text{MM} = \frac{gRT}{PV}$$

EXAMPLE #2:

Q: What is the molar mass of a gas if 0.427 grams of the gas occupies a volume of 125 mL at 20.0 °C and 0.980 atm?  
(A: 83.8 g/mole)

EXAMPLE #3:

Q: What is the density of argon gas, Ar, at a pressure of 551 torr and a temperature of 25 °C?

A: density =  $\frac{\text{mass}}{\text{volume}}$        $PV = \frac{g}{\text{MM}}RT$        $\text{MM} \cdot PV = \frac{g}{\text{MM}}RT \cdot \text{MM}$        $\frac{\text{MM} \cdot PV}{V} = \frac{gRT}{V}$

$$\frac{1}{RT} \cdot \text{MM} \cdot P = \frac{gRT}{V} \cdot \frac{1}{RT} \qquad \text{After all rearranging... Density (D) = } \frac{g}{V} = \frac{\text{MM} P}{RT}$$

$$P = \frac{551 \text{ torr}}{760 \text{ torr}} \cdot 1 \text{ atm} = 0.725 \text{ atm} \qquad T = 25 \text{ }^\circ\text{C} + 273 = 298 \text{ K}$$

$$D = \frac{(0.725 \text{ atm})(39.9 \text{ g/mole})}{(0.0821 \text{ L atm/mole K})(298 \text{ K})} = 1.18 \text{ g/L}$$

EXAMPLE #4:

Q: The density of a gas was found to be 2.0 g/L at 1.50 atm and 27 °C. What is the molar mass of the gas?  
(A: 33 g/mole)

**Applications of Ideal Gas Equation WKSHT**

- 1.) What pressure is exerted by 1.0 mole of an ideal gas contained in a 1.0 L vessel at 0.0 °C?
- 2.) What is the density of a sample of ammonia gas, NH<sub>3</sub>, if the pressure is 0.928 atm and the temperature is 63.0 °C?
- 3.) Calculate the molar mass of a gas if 4.5 L of the gas at 785 torr and 23.5 °C has a mass of 13.5 grams.
- 4.) 0.453 moles of a gas confined to a 15.0 L container exerts a pressure of 1.24 atm on the walls of the container. What is the temperature of the gas (in °C)?
- 5.) 5.4 grams of carbon dioxide are confined to a 20.0 L container at a temperature of 32.5 °C. What pressure does the gas exert?
- 6.) 2.125 grams of a gas in a 1.25 L container exert a pressure of 0.838 atm at 40.0 °C. What is the molar mass of the gas?
- 7.) To what temperature must 10.0 grams of NH<sub>3</sub> have to be heated in a 15.0 L container in order for it to exert a pressure of 3.50 atm?
- 8.) 2.0 x 10<sup>-5</sup> grams of hydrogen gas at 155 °C exert a pressure of 322.5 torr on the walls of a small cylindrical tube. What is the volume of the tube?

Answers: 1.) 22 atm; 2.) 0.572 g/L; 3.) 71 g/mole; 4.) 227 °C; 5.) 0.15 atm; 6.) 52.1 g/mole; 7.) 1090 K; 8.) 8.3 x 10<sup>-4</sup> L

**GAS LAWS NOTES**

\* Unlike Ideal Gas Equation, the "Gas Laws" describe one gas undergoing a change in conditions. The Gas Laws are also different from the Ideal Gas Equation because you do not have to convert any units except temperature that has to be in Kelvins.

Combined Gas Law:

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

\* All of the other gas laws can be derived from the combined gas law. \*

~ Boyle's Law

- describes relationship between pressure & volume when temperature is constant
- because temperature is constant, it can be excluded from the equation
- so, equation for Boyle's Law is \_\_\_\_\_
- pressure & volume are \_\_\_\_\_ proportional
- graph of pressure vs. volume would have the general shape of

EXAMPLE: A sample of gas occupies 15 liters under 2.1 atm of pressure. What would the volume of the gas be if the pressure were decreased to 1.2 atm? (Assume that temperature is constant.)

~ Charles' Law

- describes relationship between volume & temperature when pressure is constant
- because pressure is constant, it can be excluded from the equation
- so, equation for Charles' Law is \_\_\_\_\_

- volume & temperature are \_\_\_\_\_ proportional
- graph of volume vs. temperature would have the general shape of

EXAMPLE: When I purchase a helium balloon at the store (where the temperature is 25 °C) for my friend's birthday, the clerk fills the balloon to a volume of 20.0 liters. When I go outside, the balloon shrinks to a volume of 17.9 liters. What is the temperature outside?

~ Gay-Lussac's Law

- describes relationship between pressure & temperature when volume is constant
- because volume is constant, it can be excluded from the equation
- so, equation for Gay-Lussac's Law is \_\_\_\_\_
- pressure & temperature are \_\_\_\_\_ proportional
- graph of pressure vs. temperature would have the general shape of

EXAMPLE: An aerosol can has an internal pressure of 2.75 atm at room temperature (25 °C). What is the pressure in the can if I leave it outside in the sun and the temperature goes up to 35 °C?

**Gas Law Problems WKSHT**

- 1.) The gas pressure in an aerosol can is 1.5 atm at 25 °C. Assuming that the gas inside obeys the ideal gas equation, what would the pressure be if the can were heated to 450 °C?
- 2.) A pocket of gas is discovered in a deep drilling operation. The gas has a temperature of 480 °C and is at a pressure of 12.8 atm. Assume ideal behavior. What volume of the gas is required to provide 18.0 L at the surface at 1.00 atm and 22 °C?
- 3.) A fixed quantity of gas is compressed at constant temperature from a volume of 368 mL to 108 mL. If the initial pressure was 5.22 atm, what is the final pressure?
- 4.) A gas originally at 15 °C and having a volume of 182 mL is reduced in volume to 82.0 mL while its pressure is held constant. What is its final temperature?
- 5.) At 36 °C and 1.00 atm pressure, a gas occupies a volume of 0.600 L. How many liters will it occupy at 0.0 °C and 0.205 atm?
- 6.) What is the temperature at which  $9.87 \times 10^{-2}$  moles occupies 164 mL at 0.645 atm?
- 7.) Chlorine is widely used to purify municipal water supplies and to treat swimming pool waters. Suppose that the volume of a particular sample of  $\text{Cl}_2$  is 6.18 L at 0.90 atm and 33 °C. What volume will the  $\text{Cl}_2$  occupy at 107 °C and 0.75 atm?
- 8.) A gas exerts a pressure of 1.5 atm at 27 °C. The temperature is increased to 108 °C with no volume change. What is the gas pressure at the higher temperature?

Answers: 1.) 3.6 atm; 2.) 3.59 L or 3.6 L; 3.) 17.8 atm; 4.) 130 K or 130. K; 5.) 2.59 L or 2.6 L; 6.) 13.1 K  
7.) 9.2 L; 8.) 1.9 atm

**GAS STOICHIOMETRY NOTES**

- \* chemical reaction is happening
- \* deals with two different substances (at least 1 is a gas)
- \* given chemical equation
- \* assume reaction occurs at STP unless otherwise noted

**To solve stoichiometry problems... ALWAYS!!!!!!!!!!!!**

**\*\* WRITE THE BALANCED EQN & GIVEN INFORMATION! \*\***

**1.) Find moles of given element or compound.**

\* Use molar mass of given substance, if problem gives you grams.

~ or ~

\* Use 22.4 L = 1 mole of gas at STP, if problem gives you liters.

~ or ~

\* Use  $6.022 \times 10^{23}$  atoms (or molecules) = 1 mole, if problem gives you atoms/molecules.

**2.) Use mole ratio (coefficients) from balanced equation. (Same as before!)**

**3.) Find answer.**

\* Use molar mass of unknown substance, if question asks for grams.

~ or ~

\* Use 22.4 L = 1 mole of gas at STP, if problem asks for liters.

~ or ~

\* Use  $6.022 \times 10^{23}$  atoms (or molecules) = 1 mole, if problem asks for atoms/molecules.

Example # 1

How many Liters of carbon dioxide gas can be produced from the decomposition of 4.50 grams of sodium carbonate?  
 $\text{Na}_2\text{CO}_3 \rightarrow \text{Na}_2\text{O} + \text{CO}_2$  (Note: Equation is already balanced.)

$$\frac{4.50 \text{ g Na}_2\text{CO}_3}{106 \text{ g Na}_2\text{CO}_3} \times \frac{1 \text{ mole Na}_2\text{CO}_3}{1 \text{ mole Na}_2\text{CO}_3} = 0.0425 \text{ moles Na}_2\text{CO}_3$$

$$\frac{0.0425 \text{ moles Na}_2\text{CO}_3}{1} = \frac{x \text{ moles CO}_2}{1} \quad x = 0.0425 \text{ moles CO}_2$$

$$\frac{0.0425 \text{ moles CO}_2}{1} \times \frac{22.4 \text{ L CO}_2}{1 \text{ mole CO}_2} = 0.951 \text{ L CO}_2$$

Example # 2

How many liters of H<sub>2</sub> are needed to react completely with 15.0 L of N<sub>2</sub>?  
 $\text{N}_2 + \text{H}_2 \rightarrow \text{NH}_3$

Example # 3

How many grams of aluminum are needed to completely react with 16.0 L of oxygen?  
 $\text{Al} + \text{O}_2 \rightarrow \text{Al}_2\text{O}_3$  (Ans: 25.7 g)

**GAS STOICHIOMETRY (standard conditions) WKSHT.**

**\*\* assume all reactions in this section occur at STP \*\***

- How many liters of oxygen can be formed from the decomposition of 2.00 grams of  $\text{KClO}_3$ .  

$$\underline{\quad} \text{KClO}_3 \rightarrow \underline{\quad} \text{KCl} + \underline{\quad} \text{O}_2$$
- How many grams of  $\text{CaCO}_3$  are required to produce 6.00 L of  $\text{CO}_2$ ?  

$$\underline{\quad} \text{CaCO}_3 \rightarrow \underline{\quad} \text{CaO} + \underline{\quad} \text{CO}_2$$
- Determine the volume of hydrogen gas produced when 0.250 moles of zinc react with excess HCl.  

$$\underline{\quad} \text{Zn} + \underline{\quad} \text{HCl} \rightarrow \underline{\quad} \text{ZnCl}_2 + \underline{\quad} \text{H}_2$$
- How many liters of nitrogen are required to combine with 3.0 L of hydrogen in the following reaction:  

$$\underline{\quad} \text{N}_2 + \underline{\quad} \text{H}_2 \rightarrow \underline{\quad} \text{NH}_3$$
- How many liters of oxygen are needed to combine with 7.0 liters of propane in the following reaction:  

$$\underline{\quad} \text{C}_3\text{H}_8 + \underline{\quad} \text{O}_2 \rightarrow \underline{\quad} \text{CO}_2 + \underline{\quad} \text{H}_2\text{O}$$
- From the following reaction:  $\underline{\quad} \text{CH}_4 + \underline{\quad} \text{O}_2 \rightarrow \underline{\quad} \text{CO}_2 + \underline{\quad} \text{H}_2\text{O}$   
 How many liters of  $\text{CO}_2$  are formed from 32.0 grams of  $\text{CH}_4$ ?
- How many grams of Na are needed to produce 5.0 L of hydrogen?  

$$\underline{\quad} \text{Na} + \underline{\quad} \text{H}_2\text{O} \rightarrow \underline{\quad} \text{NaOH} + \underline{\quad} \text{H}_2$$
- Determine the volume of  $\text{CO}_2$  produced from burning 0.750 moles of C.  

$$\underline{\quad} \text{C} + \underline{\quad} \text{O}_2 \rightarrow \underline{\quad} \text{CO}_2$$

Answers: 1.) 0.551 L; 2.) 26.8 g; 3.) 5.60 L; 4.) 1.0 L; 5.) 35 L; 6.) 44.8 L; 7.) 10. g; 8.) 16.8 L

**DALTON'S LAW OF PARTIAL PRESSURES NOTES**

\* DEALS WITH A MIXTURE OF DIFFERENT GASES \*

- The sum of the pressures of the individual gases equals the total pressure exerted by the mixture of gases.

$$P_{\text{TOTAL}} = P_{\text{gas1}} + P_{\text{gas2}} + \dots$$

EXAMPLE: A mixture of oxygen and nitrogen exerts 1.1 atm of pressure. What is oxygen's partial pressure if the pressure of the nitrogen gas is 0.8 atm?

- The pressure of a gas "collected over water" is equal to the atmospheric pressure minus the vapor pressure of the water.

$$P_{\text{gas}} = P_{\text{atm}} - P_{\text{water}}$$

EXAMPLE: A 44.6 mL sample of carbon dioxide is collected over water at 765 mm Hg pressure and 25 °C. What is the vapor pressure of the "dry" gas? (The vapor pressure of water at 25 °C is 23.76 mm Hg. You will have to look up this value on a table or this value must be given to you.)

- The partial pressure of a gas is equal to the "mole fraction" multiplied by the total pressure.

$$P_x = \frac{(\text{moles } x)}{(\text{total moles})} \cdot P_{\text{total}} \quad \text{"}P_x\text{" = partial pressure of certain gas}$$

$$\text{"mole fraction"} = \frac{(\text{moles } x)}{(\text{total moles})}$$

EXAMPLE: A mixture of gases contains 2.0 moles of He and 4.0 moles of oxygen. If the mixture exerts a pressure of 801 torr, what is the partial pressure of the oxygen?

**GRAHAM'S LAW OF EFFUSION NOTES**

- \* compares the rates of effusion of different gases
- \* lighter gases (lower molar masses) effuse faster than heavier gases (higher molar masses)

$$\frac{\text{rate A}}{\text{rate B}} = \sqrt{\frac{\text{MM B}}{\text{MM A}}}$$

the rate of gas A compared to the rate of gas B is equal to the square root of the inverse of the molar masses of the gases

EXAMPLE 1: Compare the rates of effusion for oxygen gas and hydrogen gas.

EXAMPLE 2: An unknown gas effuses 1.18 times faster than SO<sub>2</sub>. What is the molar mass of the unknown gas?

**DALTON'S LAW & GRAHAM'S LAW WKSHT.**

- 1.) Determine the partial pressure of each gas in a container with 2.0 moles of N<sub>2</sub>, 3.0 moles of O<sub>2</sub>, and 7.0 moles of H<sub>2</sub> that has a total pressure of 850 mm Hg. (You will have 3 separate answers for this question.)
- 2.) A mixture of nitrogen and oxygen has a total pressure of 730 mm Hg. If the nitrogen has a partial pressure of 420 mm Hg, find the pressure of the oxygen.
- 3.) At an altitude of 30,000 ft., the total air pressure is only about 450. mm Hg. If the air is 21.0 % oxygen, what is the partial pressure of oxygen at this altitude?
- 4.) A mixture of 3 gases have the following pressures: oxygen = 355 mm Hg, helium = 468 mm Hg, & nitrogen = 560 mm Hg. Find the % of each gas in the mixture.
- 5.) Compare the rate of effusion of CH<sub>4</sub> and CO<sub>2</sub>. (Give answers to # 5, 6, & 7 to 3 SF's.)  
(Your answers for # 5, 6, & 7 should read " \_\_\_ effuses \_\_\_ times faster than \_\_\_.")
- 6.) Compare the rate of effusion of helium and nitrogen.
- 7.) How much faster does ammonia (NH<sub>3</sub>) effuse than HCl?
- 8.) An unknown gas effuses 4.0 times faster than O<sub>2</sub>. Find the molar mass of the unknown gas.

Answers: 1.) N<sub>2</sub> = 142 mm Hg, O<sub>2</sub> = 213 mm Hg, H<sub>2</sub> = 496 mm Hg      2.) 310 mm Hg  
 3.) 94.5 mm Hg      4.) O<sub>2</sub> = 25.7%, He = 33.8 %, N<sub>2</sub> = 40.5 %  
 5.) CH<sub>4</sub> effuses 1.66 times faster than CO<sub>2</sub>.      6.) He effuses 2.65 times faster than N<sub>2</sub>.  
 7.) NH<sub>3</sub> effuses 1.46 times faster than HCl      8.) 2.0 g/mole

**UNIT 10 REVIEW WORKSHEET**

- 1.) Convert the following pressure measurements to atmospheres.  
 (A) 151.98 kPa                      (B) 456 mm Hg                      (C) 912 torr
- 2.) What are the conditions for gas measurement at STP?
- 3.) The volume of a sample of methane gas measures 350. mL at 27.0 °C and 810. mm Hg. What is the volume (in liters) at -3.0 °C and 650. mm Hg pressure?
- 4.) How many grams of nitrogen gas are contained in a 32.6 liter container at 34.4 °C and 579 torr?
- 5.) A mixture of four gases in a container exerts a total pressure of 955 mm Hg. In this container, there are 4.50 moles of N<sub>2</sub>, 4.25 moles of CO<sub>2</sub>, 2.75 moles of H<sub>2</sub>, and 2.00 moles of O<sub>2</sub>. What is the partial pressure of H<sub>2</sub>?
- 6.) Compare the rates of effusion of carbon dioxide gas and carbon monoxide gas.
- 7.) An unknown gas effuses 1.37 times faster than chlorine gas. What is the molar mass of the unknown gas?
- 8.) Given the following unbalanced reaction:  
 \_\_\_ C<sub>5</sub>H<sub>12</sub> + \_\_\_ O<sub>2</sub> → \_\_\_ CO<sub>2</sub> + \_\_\_ H<sub>2</sub>O  
 How many liters of oxygen are needed to produce 45.7 liters of CO<sub>2</sub>?
- 9.) Given the unbalanced equation:  
 \_\_\_ Mg + \_\_\_ O<sub>2</sub> → \_\_\_ MgO  
 How many liters of oxygen gas are required to produce 45.8 grams of magnesium oxide?



## UNIT 10 - GASES

Law	Dalton's Law	Dalton's Law	Graham's Law
equation	$P_{\text{total}} = P_{\text{gas1}} + P_{\text{gas2}} + \dots$	$P_x = \frac{(\text{moles } x)}{(\text{total moles})} \cdot P_{\text{total}}$	$\frac{\text{rate A}}{\text{rate B}} = \sqrt{\frac{\text{MM B}}{\text{MM A}}}$
explanation	the sum of the pressures of the individual gases in a mixture equals the total pressure exerted by the mixture	amount of a gas in mixture is proportionate to the amount of its partial pressure	rate of gas A compared to the rate of gas B is equal to the square root of the inverse of their molar masses
when to use it	mixture of gases; only pressures given	mixture of gases; moles & total pressure given	when any form of the word "effusion" or "diffusion" is in the problem
specific units req'd?	any - but all values must have same unit of pressure	any - but all values must have same unit of pressure	no

Law	Molar Mass (from Ideal Gas Equation)	Density (from Ideal Gas Equation)
equation	$MM = \frac{g R T}{P V}$	$D = \frac{P MM}{R T}$
explanation	one gas at one set of conditions, asking about molar mass (no information about density given)	one gas at one set of conditions (involves density)
when to use it	when the problem gives: mass (grams), P, V, T	when density is mentioned (given or asked for in problem)
specific units req'd?	pressure = atm, volume = liters, temperature = Kelvins, molar mass = g/mole	pressure = atm, volume = liters, temperature = Kelvins, molar mass = g/mole, density = g/L

### UNIT 10 SUMMARY & PRACTICE WORKSHEET

- Convert the following temperatures.  
 (A) 104 °C to K      (B) -3 °C to K      (C) 67 K to °C      (D) 1671 K to °C
- Convert the following pressures.  
 (A) 635 torr to atm      (B) 104.2 kPa to mm Hg      (C) 1.45 atm to Pa
- A gas that effuses 1.19 times slower than nitrogen is added to light bulbs. What is the molecular mass of this unknown gas?
- (A) What is the molecular mass of a 0.2500 g sample of a gas at 99.8°C and 0.9131 atm in a 100.0 cm<sup>3</sup> container? (B) What is the gas in the container?
- A small 2.00 L fire extinguisher has an internal pressure of 506.6 kPa at 25°C. What volume of methyl bromide, the fire extinguisher's main ingredient, is needed to fill an empty fire extinguisher at standard pressure if the temperature remains constant?
- If 45.0 g of propane gas burns completely in the following reaction:  

$$\text{C}_3\text{H}_{8(g)} + 5 \text{O}_{2(g)} \rightarrow 3 \text{CO}_{2(g)} + 4 \text{H}_2\text{O}_{(g)}$$
 then how many liters of carbon dioxide gas will be released if the system is at STP?

## UNIT 10 - GASES

- Air in a closed cylinder is heated from 25°C to 36°C. If the initial pressure is 3.80 atm, what is the final pressure?
- At what temperature Celsius will 19.4 g of molecular oxygen, O<sub>2</sub>, exert a pressure of 1820 mm Hg in a 5.12 L cylinder?
- To what temperature must 32.0 ft<sup>3</sup> of a gas at 2.0 °C be heated for it to occupy 1.00 x 10<sup>2</sup> ft<sup>3</sup> at the same pressure? (ft<sup>3</sup> is a unit of volume)
- Determine the molar mass of a gas that has a density of 2.18 g/L at 66°C and 720 mm Hg.
- A 3.10 mL bubble of methane gas forms at the bottom of a bog where the temperature is 12°C and the pressure is 8.5 atm. The bubble rises to the surface where the temperature is 35°C and the pressure is 1.18 atm. What is the new volume of the methane bubble?
- A mixture of 2.00 moles of H<sub>2</sub>, 2.00 moles of NH<sub>3</sub>, 4.00 moles of CO<sub>2</sub> and 5.00 moles of N<sub>2</sub> exerts a total pressure of 800. torr. What is the partial pressure of the carbon dioxide gas?
- For the reaction  $2 \text{H}_{2(g)} + \text{O}_{2(g)} \rightarrow 2 \text{H}_2\text{O}_{(g)}$ , how many liters of water can be made from 5.0 L of oxygen gas and an excess of hydrogen?

ANSWERS: 1. (A) 377 K      (B) 270 K      (C) -206 °C      (D) 1398 °C  
2. (A) 0.836 atm      (B) 781.8 mm Hg      (C) 147,000 Pa      3. 39.7 g/mole  
4. (A) 83.8 g/mole      (B) Kr      5. 10.0 L      6. 68.5 L      7. 3.94 atm      8. -27 °C  
9. 859 K      10. 64.1 g/mole      11. 24 mL      12. 246 torr      13. 10. L